



[Go to Product page](#)

Datasheet for ABIN1393153 **anti-ATL1 antibody (AA 201-300) (Biotin)**

Overview

Quantity:	100 µL
Target:	ATL1
Binding Specificity:	AA 201-300
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATL1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SPG3A/Atlastin
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ATL1
Alternative Name:	SPG3A/Atlastin (ATL1 Products)

Target Details

Background:	<p>Synonyms: AD FSP, atl1, ATLA1_HUMAN, Atlastin GTPase 1, Atlastin-1, Atlastin1, Brain specific GTP binding protein, Brain-specific GTP-binding protein, FSP1, GBP-3, GBP3, GTP-binding protein 3, Guanine nucleotide-binding protein 3, Guanylate binding protein 3, hGBP3, HSN1D, Spastic paraplegia 3 protein A, SPG 3A, SPG3, SPG3A.</p> <p>Background: Atlastins are Golgi-localized, integral membrane proteins that function as GTPases. The Atlastin proteins, also designated SPG3A and guanylate-binding protein 3, comprise a Dynamin superfamily that plays a role in axonal maintenance. Hereditary spastic paraplegia (HSP) is an inherited neurodegenerative disorder that is characterized by retrograde axonal degeneration. HSP primarily affects long corticospinal neurons and causes spastic lower extremity weakness. Spastin, a microtubule (MT)-severing AAA ATPase, is a binding partner of Atlastin that is involved in membrane dynamics. This Spastin/Atlastin binding may be involved in the biochemical pathway that leads to HSP development. Mutations in the Atlastin gene (SPG3A) account for approximately 10 % of all autosomal dominant HSPs, while mutations in the Spastin gene (SPG4) account for almost 40 %.</p>
-------------	---

Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months