# antibodies .- online.com





Datasheet for ABIN1393258

## anti-ZFYVE27 antibody (AA 341-411) (Alexa Fluor 488)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	ZFYVE27	
Binding Specificity:	AA 341-411	
Reactivity:	Human, Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This ZFYVE27 antibody is conjugated to Alexa Fluor 488	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ZFYVE27
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Rabbit
Purification:	Purified by Protein A.

#### **Target Details**

Target:	ZFYVE27
Alternative Name:	ZFYVE27 (ZFYVE27 Products)

Target Details		
Background:	Synonyms: Protrudin, RP11 459F3.2, SPG33, ZFY27_HUMAN, ZFYVE27, zinc finger FYVE	
	domain containing 27, Zinc finger FYVE domain containing protein 27, Zinc finger FYVE	
	domain-containing protein 27.	
	Background: Zinc finger FYVE domain-containing protein 27 (ZFYVE27), also known as SPG33,	
	is a 411 amino acid member of the FYVE-finger family of proteins. The FYVE domain is a	
	cysteine-rich domain of about 70 amino acids that plays a role in the endosomal localization of	
	the FYVE-finger proteins, and a majority of these proteins serve as regulators of endocytic	
	membrane trafficking. ZFYVE27, a multi-pass membrane protein, is an endosomal protein that	
	binds to Spastin, a protein that is primarily involved in microtubule dynamics and severing,	
	vesicular trafficking and endosomal trafficking. Mutations in the gene encoding ZFTVE27 affect	
	neuronal intracellular trafficking in the corticospinal tract and are thought to lead to hereditary	
	spastic paraplegia (HSP), a neurodegenerative disorder, characterized by progressive paralysis	
	of the legs, which is caused by impaired axonal transport. Five isoforms of ZFYVE27 exist as a	
	result of alternative splicing events.	
Gene ID:	118813	
UniProt:	Q5T4F4	
Pathways:	Neurotrophin Signaling Pathway	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50.00	

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

50 % Glycerol.

handled by trained staff only.

ProClin

Preservative:

Precaution of Use:

### Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months