



[Go to Product page](#)

Datasheet for ABIN1393435  
**anti-Ataxin 10 antibody (AA 21-120) (Biotin)**

### Overview

Quantity:	100 µL
Target:	Ataxin 10 (ATXN10)
Binding Specificity:	AA 21-120
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ataxin 10 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ATXN10/SCA10
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse
Purification:	Purified by Protein A.

### Target Details

Target:	Ataxin 10 (ATXN10)
Alternative Name:	ATXN10/SCA10 ( <a href="#">ATXN10 Products</a> )

## Target Details

---

Background:	<p>Synonyms: Ataxin 10, Ataxin-10, ATX10_HUMAN, Atxn10, Brain protein E46 homolog, E46L, FLJ37990, HUMEEP, Like mouse brain protein E46, SCA10, Spinocerebellar ataxia 10, Spinocerebellar ataxia type 10 protein.</p> <p>Background: Spinocerebellar ataxia (SCA) is an autosomal dominant neurodegenerative disorder characterized by ataxia and selective neuronal cell loss. SCA is caused by the expansion of a translated CAG repeat, encoding a polyglutamine tract in SCA gene products, known as ataxins. The ataxin proteins are ubiquitously expressed in nervous tissue, but are primarily detected in cerebellum, brain stem and spinal cord in the central nervous system.</p> <p>Ataxin-10 is a cytoplasmic protein that belongs to the family of armadillo repeat proteins. A loss of ataxin-10 in primary neuronal cells causes increased apoptosis of cerebellar neurons. Ataxin-10 interacts with p110, an O-Linked beta-N-acetylglucosamine transferase, and may be important in the regulation of intracellular glycosylation levels and homeostasis in the brain.</p> <p>Spinocerebellar ataxia type 10 (SCA10) is an autosomal dominant disorder that causes cerebellar ataxia and seizures. SCA10 is caused by an expansion of an ATTCT pentanucleotide repeat in intron 9 of the ataxin-10 gene.</p>
-------------	---

## Application Details

---

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

## Handling

---

Expiry Date: 12 months