

[Go to Product page](#)

Datasheet for ABIN1393475

**anti-BTD antibody (AA 401-500) (Alexa Fluor 555)**

## Overview

Quantity:	100 µL
Target:	BTD
Binding Specificity:	AA 401-500
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BTD antibody is conjugated to Alexa Fluor 555
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Biotinidase
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse, Dog, Cow, Sheep, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	BTD
Alternative Name:	BTD/Biotinidase ( <a href="#">BTD Products</a> )

## Target Details

Background:	Synonyms: Biotinase, Biotinidase, Btd, BTD_HUMAN, EC 3.5.1.12.  Background: Biotin, also known as vitamin B7, is an essential water-soluble vitamin that is a cofactor in glucogenesis and in the metabolism of fatty acids and leucine. Biotinidase is a 523 amino acid enzyme that catalyzes the hydrolysis of biocytin to biotin and lysine. Secreted into extracellular space, biotinidase is expressed in liver, heart, placenta, brain, skeletal muscle, pancreas and kidney. Biotinidase contains one carbon-nitrogen hydrolase domain, which is involved in the reduction of organic nitrogen compounds and ammonia production. Defects in the gene encoding biotinidase are the cause of biotinidase deficiency, which is characterized by skin rash, ataxia, seizures, hearing loss, hypotonia and optic atrophy. These symptoms are due to the individual's inability to reutilize biotin and can, therefore, typically be treated with the addition of free biotin.
-------------	---

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months