antibodies -online.com







anti-CBLN4 antibody (AA 101-201) (Biotin)



O -	4 -	D	4	page
	$T \cap$	$Pr \cap C$	шет	nane
\cup	w	1 100	IUCL	Dauc

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	CBLN4	
Binding Specificity:	AA 101-201	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CBLN4 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CBLN4
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CBLN4
Alternative Name:	CBLN4 (CBLN4 Products)
Background:	Synonyms: CBLN 4, CBLNL1, Cerebellin 4, Cerebellin 4 precursor, Cerebellin like glycoprotein 1,

Cerebellin precursor like 1, Cerebellin4, CBLN4_HUMAN.

Background: Cerebellin (CER), which was originally isolated from rat cerebellum, is a hexadecapeptide derived from a larger precursor called Cerebellin 1, also designated precerebellin 1 or Cbln1. Four propeptides, Cerebellin 1, Cerebellin 2 (Cbln2), Cerebellin 3 (Cbln3) and Cerebellin 4 (Cbln4), comprise the precerebellin subfamily within the C1q protein family. Cerebellin family members act as transneuronal regulators of synapse development and synaptic plasticity in various brain regions. Cerebellin and its metabolite, des-Ser(1)Cer, are also expressed in several extra-cerebellar tissues, including adrenal gland. Cerebellin 1, 2 and 3 assemble into homomeric and heteromeric complexes, thereby influencing each other?s degradation and secretion. Cerebellin 3 is not able to form homomeric complexes, and can only be secreted upon forming a heteromeric complex with Cerebellin 1. Decreased concentrations of Cerebellin have been found in the brain of patients with olivopontocerebellar atrophy (OPCA) and Shy-Drager syndrome, suggesting a role for Cerebellin in the pathology of these diseases.

Application Details

Expiry Date:

WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500
For Research Use only
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Freservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20 °C for 12 months.

12 months