antibodies -online.com





anti-CSAD antibody (AA 401-493) (Biotin)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | CSAD | |
| Binding Specificity: | AA 401-493 | |
| Reactivity: | Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This CSAD antibody is conjugated to Biotin | |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CSAD |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Predicted Reactivity: | Human,Dog,Horse |
| Purification: | Purified by Protein A. |

Target Details

| Target: | CSAD |
|-------------------|----------------------|
| Alternative Name: | CSAD (CSAD Products) |

Target Details

| Background: | Synonyms: CSAD, CSAD_HUMAN, CSD, Cysteine sulfinic acid decarboxylase, Cysteine-sulfinate | |
|---------------------|---|--|
| | decarboxylase, Sulfinoalanine decarboxylase. | |
| | Background: CSAD is a 493 amino acid protein that exists as a homodimer and belongs to the | |
| | group II decarboxylase family. CSAD catalyzes the conversion of 3-sulfino-L-alanine to | |
| | hypotaurine and carbon dioxide, binds pyridoxal phosphate as a cofactor and undergoes | |
| | alternative splicing to produce three isoforms. The gene encoding CSAD maps to human | |
| | chromosome 12, which encodes over 1,100 genes and comprises approximately 4.5 $\%$ of the | |
| | human genome. Chromosome 12 is associated with a variety of diseases and afflictions, | |
| | including hypochondrogenesis, achondrogenesis, Kniest dysplasia, Noonan syndrome and | |
| | trisomy 12p, which causes facial developmental defects and seizure disorders. | |
| Gene ID: | 51380 | |
| Application Details | | |
| Application Notes: | WB 1:300-5000 | |
| | IHC-P 1:200-400 | |
| | IHC-F 1:100-500 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C for 12 months. | |
| Expiry Date: | 12 months | |