# antibodies -online.com





## anti-DCDC2 antibody (AA 85-168) (Alexa Fluor 350)



Go to Product page

| ( ) | ve     | K\ / |   | A . |
|-----|--------|------|---|-----|
|     | $\cup$ | 1 V/ | - | V۷  |
|     |        |      |   |     |

| Quantity:            | 100 μL   |  |
|----------------------|--|--|
| Target:              | DCDC2  |  |
| Binding Specificity: | AA 85-168  |  |
| Reactivity:          | Human  |  |
| Host:                | Rabbit   |  |
| Clonality:           | Polyclonal   |  |
| Conjugate:           | This DCDC2 antibody is conjugated to Alexa Fluor 350   |  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human DCDC2 |
|-----------------------|---|
| Isotype:              | IgG   |
| Predicted Reactivity: | Human,Mouse,Rat,Cow,Sheep,Horse                           |
| Purification:         | Purified by Protein A.                                    |

### **Target Details**

| Target:   | DCDC2                  |
|---|------------------------|
| Alternative Name:   | DCDC2 (DCDC2 Products) |
| Background: Synonyms: RU2, RU2S, DCDC2A, DFNB66, NPHP19, Doublecortin domain-containing pro |                        |

#### **Target Details**

| rarget Details      |   |
|---------------------|---|
|                     | Protein RU2S, DCDC2, KIAA1154   |
|                     | Background: Protein that plays a role in the inhibition of canonical Wnt signaling pathway. May |
|                     | be involved in neuronal migration during development of the cerebral neocortex. Involved in the |
|                     | control of ciliogenesis and ciliary length (PubMed:25601850).                                   |
| Gene ID:            | 51473   |
| UniProt:            | Q9UHG0  |
| Application Details |   |
| Application Notes:  | IF(IHC-P) 1:50-200  |
|                     | IF(IHC-F) 1:50-200  |
|                     | IF(ICC) 1:50-200  |
| Restrictions:       | For Research Use only   |
| Handling            |   |
| Format:             | Liquid  |
| Concentration:      | 1 μg/μL   |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and     |
|                     | 50 % Glycerol.  |
| Preservative:       | ProClin   |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be             |
|                     | handled by trained staff only.  |
| Storage:            | -20 °C  |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.               |
| Expiry Date:        | 12 months   |