antibodies .- online.com





anti-Fam53a antibody (AA 255-340) (Alexa Fluor 555)



| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | Fam53a | |
| Binding Specificity: | AA 255-340 | |
| Reactivity: | Human, Dog | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This Fam53a antibody is conjugated to Alexa Fluor 555 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human DNTNP/FAM53A |
|-----------------------|--|
| In the second | |
| Isotype: | IgG |
| Cross-Reactivity: | Dog, Human |
| Predicted Reactivity: | Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | Fam53a |
|-------------------|--------------------------------|
| Alternative Name: | DNTNP/FAM53A (Fam53a Products) |

Target Details

| rai got botano | | |
|---------------------|---|--|
| Background: | Synonyms: DNTNP, Dorsal neural-tube nuclear protein, FA53A_HUMAN, FAM53A, family with | |
| | sequence similarity 53, member A, Protein FAM53A. | |
| | Background: Dorsal neural-tube nuclear protein is a 398 amino acid nuclear protein. DNTNP is | |
| | thought to play an important role in neural development by specifying dorsal cell fates within | |
| | the neural tube. Widely expressed in the dorsal neural tube, DNTNP is most highly expressed in | |
| | the dorsal regions of the midbrain, the hindbrain, the diencephalon, and the spinal neural tube | |
| | and is expressed at lower levels in the branchial arches, the telencephalon, the heart, and | |
| | somites of developing embryos. DNTNP is encoded by a gene located on chromosome 4 which | |
| | contains many genes including the Huntingtin gene, which is found to encode an expanded | |
| | glutamine tract in cases of Huntington's disease. | |
| Pathways: | Protein targeting to Nucleus | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| | | |

12 months

Expiry Date: