antibodies.com

Datasheet for ABIN1393749 anti-JARID2 antibody (AA 301-400) (Alexa Fluor 350)



Overview

Quantity:	100 µL
Target:	JARID2
Binding Specificity:	AA 301-400
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This JARID2 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Jarid2/Jumonji
lsotype:	lgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.
T . D . N	
Target Details	

Target:	JARID2
Alternative Name:	Jarid2/Jumonji (JARID2 Products)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1393749 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Deelvaround	Quanuma: IMI Dratain lumanii lumanii/ADD daraain aastaising sestaising (ADD)
Background:	Synonyms: JMJ, Protein Jumonji, Jumonji/ARID domain-containing protein 2, JARID2 Background: Regulator of histone methyltransferase complexes that plays an essential role in
	embryonic development, including heart and liver development, neural tube fusion process and
	hematopoiesis. Acts by modulating histone methyltransferase activity and promoting the
	recruitment of histone methyltransferase complexes to their target genes. Binds DNA and
	mediates the recruitment of the PRC2 complex to target genes in embryonic stem cells. Does
	not have histone demethylase activity but regulates activity of various histone methyltransferase complexes. In embryonic stem cells, it associates with the PRC2 complex
	and inhibits trimethylation of 'Lys-27' of histone H3 (H3K27me3) by the PRC2 complex, thereby
	playing a key role in differentiation of embryonic stem cells and normal development. In cardiac
	cells, it is required to repress expression of cyclin-D1 (CCND1) by activating methylation of 'Lys
	9' of histone H3 (H3K9me) by the GLP1/EHMT1 and G9a/EHMT2 histone methyltransferases.
	Also acts as a transcriptional repressor of ANF via its interaction with GATA4 and NKX2-5.
	Participates in the negative regulation of cell proliferation signaling.
Gene ID:	3720
UniProt:	Q92833
Pathways:	Chromatin Binding
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1393749 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Handling	
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months