
# antibodies -online.com







# anti-KIAA0319 antibody (AA 682-760) (Alexa Fluor 350)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	KIAA0319	
Binding Specificity:	AA 682-760	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KIAA0319 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human DYX2/KIAA0319	
Isotype:	IgG	
Predicted Reactivity: Human,Mouse,Rat,Cow,Sheep,Pig,Horse		
Purification:	Purified by Protein A.	

## **Target Details**

Target:	KIAA0319
Alternative Name: DYX2/KIAA0319 (KIAA0319 Products)	
Background: Synonyms: DLX 2, DLX2, DYLX 2, DYLX2, Dyslexia susceptibility 2, Dyslexia-associated p	

KIAA0319 DYX 2, DYX2, K0319\_HUMAN, Kiaa0319, MGC176717.

Background: DYX2 is a 1072 amino acid single-pass transmembrane protein that contains one MANSC domain and two PKD (Polycystic Kidney Disease) domains, which are usually found in the extracellular regions of proteins and are involved in protein-protein interactions. In DYX2, it is likely that its PKD domains mediate the interaction between neurons and glial fibers during neuronal migration. When overexpressed, this plasma membrane protein colocalizes with EEA1 (early endosome antigen 1) in large intracellular vesicles, suggesting that it is endocytosed and recycled. DYX2 is highly expressed in brain cortex, cerebellum, amygdala, putamen and hippocampus. Defects in the gene encoding DYX2 may be the cause of dyslexia type 2, a relatively common disorder that is characterized by reading performance impairment in the absence of sensory or neurologic disability. There are three isoforms of DYX2 that are produced as a result of alternative splicing events

### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	