antibodies -online.com





anti-KIAA0319 antibody (AA 682-760) (Alexa Fluor 555)



Go to Product page

\sim			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	KIAA0319	
Binding Specificity:	AA 682-760	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KIAA0319 antibody is conjugated to Alexa Fluor 555	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DYX2/KIAA0319	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

Target Details

Target:	KIAA0319 DYX2/KIAA0319 (KIAA0319 Products)	
Alternative Name:		
Background:	Synonyms: DLX 2, DLX2, DYLX 2, DYLX2, Dyslexia susceptibility 2, Dyslexia-associated protein	

KIAA0319 DYX 2, DYX2, K0319_HUMAN, Kiaa0319, MGC176717.

Background: DYX2 is a 1072 amino acid single-pass transmembrane protein that contains one MANSC domain and two PKD (Polycystic Kidney Disease) domains, which are usually found in the extracellular regions of proteins and are involved in protein-protein interactions. In DYX2, it is likely that its PKD domains mediate the interaction between neurons and glial fibers during neuronal migration. When overexpressed, this plasma membrane protein colocalizes with EEA1 (early endosome antigen 1) in large intracellular vesicles, suggesting that it is endocytosed and recycled. DYX2 is highly expressed in brain cortex, cerebellum, amygdala, putamen and hippocampus. Defects in the gene encoding DYX2 may be the cause of dyslexia type 2, a relatively common disorder that is characterized by reading performance impairment in the absence of sensory or neurologic disability. There are three isoforms of DYX2 that are produced as a result of alternative splicing events

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Application Details

Precaution of Use:

Storage Comment:

Storage:

Expiry Date:

Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
5.66	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Buffer:		
Buffer: Preservative:		

handled by trained staff only.

-20 °C

12 months