antibodies -online.com





anti-LPPR4 antibody (AA 251-360) (Alexa Fluor 350)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | LPPR4 | |
| Binding Specificity: | AA 251-360 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This LPPR4 antibody is conjugated to Alexa Fluor 350 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human LPPR4 |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Chicken,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | LPPR4 | |
|--|------------------------|--|
| Alternative Name: | LPPR4 (LPPR4 Products) | |
| Background: Synonyms: Brain-specic phosphatidic acid phosphatase-like protein 1, KIAA0455, Lip | | |

phosphate phosphatase-related protein type 4, Lppr4, LPPR4_HUMAN, Plasticity-related gene 1 protein, PRG-1.

Background: Phosphatidate phosphatases are a family of integral membrane glycoproteins that dephosphorylate a variety of lipid phosphates and play a role in signal transduction via the phospholipase D pathway. PAP-2 proteins function independently of Mg2+ and are insensitive to NEM (N-ethylmaleimide) inhibition. The lipid phosphates degraded by this family include ceramide 1-phosphate (C1P), sphingosine 1-phosphate (S1P), phosphatidic acid (PA) and lysophosphatidic acid (LPA). LPPR4 (lipid phosphate phosphatase-related protein type 4), also known as LPR4, PHP1, PRG1 or PRG-1, is a 763 amino acid multi-pass membrane protein that belongs to the PA-phosphatase related phosphoesterase family. Exclusively expressed in neurons, LPPR4 hydrolyzes lysophosphatidic acid (LPA) and facilitates axonal outgrowth during development and regenerative sprouting. LPPR4 exists as two alternatively spliced isoforms and is encoded by a gene located on human chromosome 1p21.2.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 | |
|--------------------|--|--|
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |