



[Go to Product page](#)

Datasheet for ABIN1393854

anti-LRRTM2 antibody (AA 81-200) (Alexa Fluor 647)

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | LRRTM2 |
| Binding Specificity: | AA 81-200 |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This LRRTM2 antibody is conjugated to Alexa Fluor 647 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human LRRTM2 |
| Isotype: | IgG |
| Cross-Reactivity: | Rat |
| Predicted Reactivity: | Human, Mouse, Dog, Cow, Pig, Horse, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | LRRTM2 |
| Alternative Name: | LRRTM2 (LRRTM2 Products) |

Target Details

| | |
|-------------|---|
| Background: | <p>Synonyms: KIAA0416, Leucine rich repeat transmembrane neuronal 2, Leucine rich repeat transmembrane neuronal 2 protein, Leucine rich repeat transmembrane protein 2, Leucine-rich repeat neuronal 2 protein, Leucine-rich repeat transmembrane neuronal protein 2, LRRN2, LRRT2_HUMAN, Lrrtm2.</p> <p>Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic Alpha/Beta horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. The LRRTM protein family plays a role in the regulation of various cellular events during nervous system development. Localizing predominantly to the nervous system, LRRTM family members are known to exhibit synaptogenic activity. LRRTM2 (leucine rich repeat transmembrane neuronal 2), also known as LRRN2, is a 516 amino acid single-pass type I membrane protein involved in the development maintenance of the vertebrate nervous system. Expressed in kidney and neuronal tissues, LRRTM2 contains ten LRR repeats and belongs to the LRRTM family. LRRTM2 is encoded by a gene that maps to human chromosome 5q31.2.</p> |
|-------------|---|

Pathways: [Synaptic Membrane](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Handling

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months