antibodies -online.com





anti-LRRTM4 antibody (AA 135-250) (Alexa Fluor 488)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	LRRTM4
Binding Specificity:	AA 135-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LRRTM4 antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LRRTM4
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	LRRTM4
Alternative Name:	LRRTM4 (LRRTM4 Products)
Background:	Synonyms: Leucine rich repeat transmembrane neuronal 4, Leucine rich repeat transmembrane

neuronal 4 protein, Leucine rich repeat transmembrane neuronal protein 4, Leucine-rich repeat transmembrane neuronal 4 protein, Leucine-rich repeat transmembrane neuronal protein 4, LRRT4_HUMAN, Lrrtm4.

Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic / horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. The LRRTM protein family plays a role in the regulation of various cellular events during nervous system development. Localizing predominantly to the nervous system, LRRTM family members are known to exhibit synaptogenic activity. LRRTM4 (leucine-rich repeat transmembrane neuronal protein 4) is a 590 amino acid member of the LRRTM protein family. Expressed in neuronal tissues, LRRTM4 may play a role in the development and maintenance of the vertebrate nervous system. A single-pass type I membrane protein, LRRTM4 contains 10 LRR repeats. LRRTM4 is expressed as two isoforms produced by alternative splicing.

Pathways:

Synaptic Membrane

Application Details

Application Notes:

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months