antibodies

Datasheet for ABIN1393938 anti-MTPN antibody (AA 41-118) (Alexa Fluor 647)



Overview

| Quantity: | 100 µL |
|----------------------|--|
| Target: | MTPN |
| Binding Specificity: | AA 41-118 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MTPN antibody is conjugated to Alexa Fluor 647 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Myotrophin/GCDP |
|-----------------------|---|
| Isotype: | lgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | MTPN |
|-------------------|---|
| Alternative Name: | Myotrophin/GCDP (MTPN Products) |
| Background: | Synonyms: FLJ31098, FLJ99857, GCDP, Granule cell dferentiation protein, MTPN, |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1393938 | 03/07/2024 | Copyright antibodies-online. All rights reserved. MTPN_HUMAN, Myotrophin, Protein V 1, Protein V-1, V 1, MTPN_HUMAN.

Background: Myotrophin (V-1 protein) is a ubiquitously expressed cytoplasmic protein that can translocate to the nucleus during sustained NFkB activation. The gene encoding for this protein localizes to chromosome 7q33. Myotrophin may be involved in cerebellar morphogenesis and contains an acetylated N-terminus and 2.5 internal 33 amino acid ankyrin repeats. It is important in the differentiation of cerebellar neurons, particularly of granule cells. The 117 amino acid protein has been associated with, and able to induce, cardiac hypertrophy. Myotrophin increases protooncogene, ANF and Beta-Myosin heavy chain transcript levels. Myotrophin is upregulated when myocytes undergo cyclic stretch or are treated with tumor necrosis factor Alpha (TNF Alpha) or interleukin-1Beta. Highest levels of Myotrophin are detected in brain and lowest levels in skeletal muscle.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1393938 | 03/07/2024 | Copyright antibodies-online. All rights reserved.