# antibodies - online.com





Datasheet for ABIN1393969

## anti-NETO2 antibody (AA 151-250) (Biotin)



( )	ve	K\ /		A .
	$\cup$	1 V/	-	V۷

Quantity:	100 μL
Target:	NETO2
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NETO2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NETO2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig
Purification:	Purified by Protein A.

### **Target Details**

Target:	NETO2
Alternative Name:	NETO2 (NETO2 Products)
Background:	Synonyms: Brain specic transmembrane protein, Brain specic transmembrane protein

containing 2 CUB and 1 LDL receptor class A domains protein 2, Brain-specic transmembrane protein containing 2 CUB and 1 LDL-receptor class A domains protein 2, BTCL2, FLJ10430, FLJ14724, FLJ90456, NEOT2, NETO 2, Neto2, NETO2\_HUMAN, Neuropilin NRP and tolloid TLL like 2, Neuropilin and tolloid like protein 2, Neuropilin and tolloid-like protein 2.

Background: NETO2 is a 525 amino acid single-pass type I membrane protein that contains two CUB domains and one LDL-receptor class A domain. Expressed as multiple alternatively spliced isoforms, NETO2 is thought to play a role in the development and maintenance of neuronal circuitry, possibly playing a role in proper brain function. The gene encoding NETO2 maps to human chromosome 16, which is associated with a variety of genetic disorders, encodes over 900 genes and comprises nearly 3 % of the human genome. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is associated with chromosome 16, as is Crohn's disease, which is a gastrointestinal inflammatory condition.

#### **Application Details**

Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months