# antibodies .- online.com





# anti-ABCB9 antibody (AA 321-420) (AbBy Fluor® 555)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL
Target:	ABCB9
Binding Specificity:	AA 321-420
Reactivity:	Mouse, Rat, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ABCB9 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ABCB9
Isotype:	IgG
Cross-Reactivity:	Mouse, Pig, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Horse,Chicken
Purification:	Purified by Protein A.

### **Target Details**

Target:	ABCB9
Alternative Name:	ABCB9 (ABCB9 Products)

#### Target Details

Bac	kar	oun	d:

Synonyms: ABC transporter 9 protein, ABCB 9, ATP binding cassette sub family B MDR/TAP member 9, ATP binding cassette sub family B member 9 precursor, ATP binding cassette transporter 9, EST122234, ABCB9, KIAA1520, TAP like protein, TAPL, ABCB9\_HUMAN. Background: ATP-binding cassette (ABC) transporters are an evolutionarily conserved family of widely-expressed proteins that use ATP hydrolysis to catalyze the transport of various molecules across extracellular and intracellular membranes. As the largest family of transmembrane proteins, ABC genes comprise several subfamilies (ABC1, ABCA, ABCE, ABCF, MDR/TAP, MRP, ALD, OABP, GCN20 and White (also known as ABCG)). In bacteria, ABC transporters are used to import compunds that cannot be obtained by diffusion. Eukaryotic ABC transporters are largely responsible for trafficking hydrophobic compounds either within the cell as part of a metabolic process or outside the cell for transport to other organs, or for secretion from the body. ABCB9 (also designated Transporter associated with antigen processing (TAP)-like or TAPL) forms a homodimer, which is localized in lysosomes. It functions as an ATP-dependent peptide transporter that shows a broad peptide specificity ranging from 6-mer up to 59-mer peptides. ABCB9 transports these peptides with low affinity but high efficiency.

Pathways:

Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process

#### **Application Details**

Application Notes:	

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

## Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months