antibodies -online.com





anti-NGRN antibody (AA 96-200) (Biotin)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	NGRN	
Binding Specificity:	AA 96-200	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NGRN antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Neugrin	
Isotype:	IgG	
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Horse, Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	NGRN	
Alternative Name: Neugrin (NGRN Products)		
Background:	ckground: Synonyms: DSC92, FI58Gm, Mesenchymal stem cell protein DSC92, Neugrin, Neugrin neu	

outgrowth associated, Neurite outgrowth associated protein, Neurite outgrowth-associated protein, Ngrn, NGRN_HUMAN, Spinal cord-derived protein FI58G.

Background: Neugrin, also known as NGRN, mesenchymal stem cell protein DSC92, neurite outgrowth-associated protein or spinal cord-derived protein FI58G, is a 291 amino acid protein that plays a role in neuronal differentiation and belongs to the neugrin family. As both a secreted and nuclear protein, neugrin exists as two alternatively spliced isoforms and is highly expressed in skeletal muscle, brain and heart. Neugrin is upregulated in neuroblastostoma cells by retinoic acid treatment and is encoded by a gene that maps to human chromosome 15q26.1. Chromosome 15 houses over 700 genes and comprises nearly 3 % of the human genome. Angelman syndrome, Prader-Willi syndrome, Tay-Sachs disease and Marfan syndrome are all associated with defects in chromosome 15-localized genes.

Application Details

Application Notes:	WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months