

[Go to Product page](#)

Datasheet for ABIN1394091

anti-NEUROD2 antibody (AA 121-250) (Alexa Fluor 350)

Overview

Quantity:	100 µL
Target:	NEUROD2
Binding Specificity:	AA 121-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NEUROD2 antibody is conjugated to Alexa Fluor 350
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Neuro D2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NEUROD2
Alternative Name:	Neuro D2 (NEUROD2 Products)
Background:	Synonyms: bHLHa1, Class A basic helix-loop-helix protein 1, class A basic helix loop helix

Target Details

protein 1, NDF2_HUMAN, NDR2, NDRF, Neuro-D2, NeuroD-related factor, NeuroD2, Neurogenic dferentiation factor 2, NDF2_HUMAN, neuroD related factor, NeuroD2, neurogenic basic helix loop helix protein, neurogenic dferentiation 2, Neurogenic dferentiation factor 2, neuronal dferentiation 2.

Background: Members of the myogenic determination family are basic helix-loop-helix (bHLH) proteins that can be separated into two classes, both of which work together to activate DNA transcription. Class A proteins include the ubiquitously expressed E-box binding factors, namely E2A, ITF-2 and HEB, while class B proteins, such as MyoD, myogenin and Neuro D (BETA2), are transiently expressed and exhibit a much more limited tissue distribution. Working in opposition to these positively acting factors are a specialized group of basic helix-loop-helix (bHLH) transcription factors that function as dominant negative regulators and are involved in cell lineage determination and differentiation. Neuro D2 (neurogenic differentiation 2), also known as NDRF, NEUROD2 or bHLHa1, is a 382 amino acid nuclear protein that contains one bHLH domain and functions to induce neurogenic differentiation, playing an important role in the maintenance and determination of cell fate.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months