antibodies .- online.com





Datasheet for ABIN1394142

anti-PNMA1 antibody (AA 165-270) (Alexa Fluor 647)

Go to Product page

| () | 11/0 | K\ / | iew | 1 |
|-----|--------|--------|---------------------|---|
| | \cup | 'I V/I | $\square \vee \vee$ | ı |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|---|--|
| Target: | PNMA1 | |
| Binding Specificity: | AA 165-270 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This PNMA1 antibody is conjugated to Alexa Fluor 647 | |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human PNMA1 | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | PNMA1 | |
|-------------------|---|--|
| Alternative Name: | PNMA1 (PNMA1 Products) | |
| Background: | Synonyms: 37 kDa neuronal protein, MA1, Paraneoplastic antigen MA1, | |

Background: Paraneoplastic neurological disorders (PNDs) are rare syndromes that are caused by, or associated with, an underlying neoplasm. The most common neoplasm among young male patients is testicular cancer, but the leading cause among other patients is lung cancer. Most PNDs are caused by an immune response against onconeural antigens, causing progressive neurological damage. The paraneoplastic antigen MA family contains three known members: MA1, MA2 and MA3. MA1, also designated neuron- and testis-specific protein 1, is a nucleolar protein in normal cells but localizes to the cytoplasm of tumor cells. MA2, also designated onconeuronal antigen MA2, is a nucleolar protein expressed in brain and testis. MA3 is highly expressed in brain and testis and is expressed at low levels in heart, trachea and kidney.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |