# antibodies -online.com







## anti-RILPL2 antibody (AA 111-211) (Alexa Fluor 350)



$\sim$						
	1//	Д	r۱	1	<b>Θ</b> 1	٨

Quantity:	100 μL	
Target:	RILPL2	
Binding Specificity:	AA 111-211	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This RILPL2 antibody is conjugated to Alexa Fluor 350	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human RILPL2	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	RILPL2	
Alternative Name:	RILPL2 (RILPL2 Products)	

#### **Target Details**

#### Background:

Synonyms: FLJ30380, FLJ32372, MGC7036, p40phox-binding protein, Rab-interacting lysosomal protein-like 2, RILP-like protein 2, RILPL2, RIPL2\_HUMAN, RLP2, Rab interacting lysosomal protein-like 2.

Background: RILPL2 is a 211 amino acid protein that belongs to the RILPL family. RILPL2 does not regulate lysosomal morphology or distribution. RILPL2 shares 32 % and 18 % amino acid identity with RILPL1 and RILP, respectively. RILPL2 as a novel interacting partner for the actin-based molecular motor MyoVa, and has a novel role for RILPL2 in controlling neuronal morphogenesis. It has been suggested that there is also a novel role for RILPL2 in the regulation of cellular shape and dendritic-spine morphogenesis, probably via the Rac1-Pak pathway. PCR analysis of human tissues detects highest RILPL2 expression in lung, followed by placenta. Lower expression is detected in liver, kidney, pancreas, heart and brain, but no expression is detected in skeletal muscle. The RILPL2 gene is conserved in chimpanzee, dog, cow, mouse, rat, chicken and zebrafish, and maps to human chromosome 12q24.31.

#### **Application Details**

Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	