antibodies -online.com







anti-SATB2 antibody (AA 451-485) (Alexa Fluor 350)



Go to Product page

()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	SATB2	
Binding Specificity:	AA 451-485	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SATB2 antibody is conjugated to Alexa Fluor 350	
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SATB2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	SATB2
Alternative Name:	SATB2 (SATB2 Products)

Target Details

Background:	Synonyms: GLSS, DNA-binding protein SATB2, Special AT-rich sequence-binding protein 2,	
	SATB2, KIAA1034	
	Background: Binds to DNA, at nuclear matrix- or scaffold-associated regions. Thought to	
	recognize the sugar-phosphate structure of double-stranded DNA. Transcription factor	
	controlling nuclear gene expression, by binding to matrix attachment regions (MARs) of DNA	
	and inducing a local chromatin-loop remodeling. Acts as a docking site for several chromatin	
	remodeling enzymes and also by recruiting corepressors (HDACs) or coactivators (HATs)	
	directly to promoters and enhancers. Required for the initiation of the upper-layer neurons (UL1)	
	specific genetic program and for the inactivation of deep-layer neurons (DL) and UL2 specific	
	genes, probably by modulating BCL11B expression. Repressor of Ctip2 and regulatory	
	determinant of corticocortical connections in the developing cerebral cortex. May play an	
	important role in palate formation. Acts as a molecular node in a transcriptional network	
	regulating skeletal development and osteoblast differentiation.	
Gene ID:	23314	
UniProt:	Q9UPW6	
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months