

[Go to Product page](#)

Datasheet for ABIN1394349

**anti-SORCS2 antibody (AA 201-300) (Alexa Fluor 350)**

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | SORCS2   |
| Binding Specificity: | AA 201-300   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This SORCS2 antibody is conjugated to Alexa Fluor 350  |
| Application:         | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human SORCS2 |
| Isotype:              | IgG  |
| Cross-Reactivity:     | Human  |
| Predicted Reactivity: | Mouse,Rat  |
| Purification:         | Purified by Protein A.                                     |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | SORCS2                                     |
| Alternative Name: | SORCS2 ( <a href="#">SORCS2 Products</a> ) |

## Target Details

|             |  |
|-------------|--|
| Background: | <p>Synonyms: KIAA1329, SORCS 2, Sortilin related VPS10 domain containing receptor 2, VPS10 domain receptor protein, VPS10 domain receptor protein SORCS 2, SORC2_HUMAN.</p> <p>Background: There are three SorCS genes that have diverse, partially overlapping functions in the central nervous system. In the developing and mature central nervous system, SorCS1, SorCS2 and SorCS3 genes are expressed in a combinatorial, non-overlapping pattern. SorCS proteins show homology to the mosaic receptor SorLA and the neurotensin receptor sortilin, based on a common VPS10 domain, which is the hallmark of the SorCS receptor family.</p> <p>SorCS2 (sortilin-related VPS10 domain containing receptor 2) is a 1,150 amino acid single-pass type I membrane protein that is highly expressed in brain and kidney. Containing six BNR repeats and a single PKD domain, SorCS2 is encoded by a gene that maps to human chromosome 4, which encodes nearly 6 % of the human genome and has the largest gene deserts (regions of the genome with no protein encoding genes) of all of the human chromosomes.</p> |
|-------------|--|

## Application Details

|                    |   |
|--------------------|---|
| Application Notes: | <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p> |
| Restrictions:      | For Research Use only   |

## Handling

|                    |  |
|--------------------|--|
| Format:            | Liquid   |
| Concentration:     | 1 µg/µL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |