antibodies -online.com





anti-NUMBL antibody (AA 75-120) (Biotin)



Go to Product page

| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL |
|----------------------|---|
| Target: | NUMBL |
| Binding Specificity: | AA 75-120 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NUMBL antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human NUMBL |
|-----------------------|---|
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | NUMBL | |
|-------------------|--|--|
| Alternative Name: | NUMBL (NUMBL Products) | |
| Background: | Synonyms: CAG 3A, CAG3A, CTG 3a, CTG3a, NBL, NUMB Drosophilia Homolog Like, Numb | |

homolog Drosophila like, Numb homolog like, Numb like protein, NUMB R, Numb-like protein, Numb-R, Numb-related protein, NUMBL, NUMBL_HUMAN, NUMBR, TNRC 23, TNRC23. Background: In Drosophila, neuronal cell fate decisions are directed by NUMB, a signaling adapter protein with two protein-protein interaction domains, namely a phosphotyrosine-binding domain and a proline-rich SH3-binding region (PRR). The mammalian NUMB homolog plays a role in the determination of cell fate during development and binds with a variety of proteins, including Eps15, LNX1 and Notch 1. NumbL (NUMB-like protein), also known as Numb-R, NBL, CAG3A, CTG3a, NUMBLIKE or TNRC23, is a 609 amino acid cytoplasmic protein that, like NUMB, is thought to play a role in cell fate. Expressed at high levels in developing brain tissue, NumbL contains one PID (phosphotyrosine interaction domain) and plays an important role in neuronal differentiation, possibly associating with Eps15 and Notch 1. In mice, deletion of the NumbL gene is associated with early embryonic death, suggesting an essential role for NumbL in early development.

Pathways:

Cell-Cell Junction Organization

Application Details

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Expiry Date: | 12 months |