



[Go to Product page](#)

Datasheet for ABIN1394519

anti-PPP2R5C antibody (AA 390-435) (Alexa Fluor 555)

Overview

Quantity:	100 µL
Target:	PPP2R5C
Binding Specificity:	AA 390-435
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PPP2R5C antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PPP2R5C
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PPP2R5C
Alternative Name:	PPP2R5C (PPP2R5C Products)

Target Details

Background:	<p>Synonyms: 2A5G_HUMAN, B' alpha regulatory subunit, B56G, KIAA0044, MGC23064, PP2A B subunit B' gamma isoform, PP2A B subunit B56 gamma isoform, PP2A B subunit isoform B"-gamma, PP2A B subunit isoform B56-gamma, PP2A B subunit isoform PR61-gamma, PP2A B subunit isoform R5-gamma, PP2A B subunit PR61 gamma isoform, PP2A B subunit R5 gamma isoform, Ppp2r5c, PR61G, Protein phosphatase 2 regulatory subunit B B56 gamma isoform, Protein phosphatase 2 regulatory subunit B' gamma, Renal carcinoma antigen NY REN 29, Renal carcinoma antigen NY-REN-29, Serine/threonine protein phosphatase 2A 56 kDa regulatory subunit gamma isoform, Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit gamma isoform, 2A5G_HUMAN.</p> <p>Background: In eukaryotes, the phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions, including division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the protein phosphatases. In general, the protein phosphatase (PP) holoenzyme is a trimeric complex composed of a regulatory subunit, a variable subunit, and a catalytic subunit. Four major families of protein phosphatase catalytic subunits have been identified, designated PP1, PP2A, PP2B (calcineurin) and PP2C. An additional protein phosphatase catalytic subunit, PPX (also known as PP4) is a putative member of a novel PP family. The PP2A family comprises subfamily members PP2A Alpha and PP2A Beta. The PP2A catalytic subunit associates with a variety of regulatory subunits. Regulatory subunits include PP2A-A-Alpha and -A-Beta, PP2A-B-Alpha and -B-Beta, PP2A-C-Alpha and -C-Beta, PP2A-B56-Alpha, -B56-Beta, -B56-gamma and -B56-Delta.</p>
-------------	---

Pathways:	PI3K-Akt Signaling
-----------	------------------------------------

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and

Handling

50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months