

[Go to Product page](#)

Datasheet for ABIN1394616

**anti-GRIK3 antibody (AA 501-600) (Alexa Fluor 647)**

## Overview

Quantity:	100 µL
Target:	GRIK3
Binding Specificity:	AA 501-600
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GRIK3 antibody is conjugated to Alexa Fluor 647
Application:	Western Blotting (WB), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GRIK3/GLR7
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

## Target Details

Target:	GRIK3
Alternative Name:	GRIK3/GLR7 ( <a href="#">GRIK3 Products</a> )

## Target Details

Background:	<p>Synonyms: EAA5, Excitatory amino acid receptor 5, GLR 7, GLR7, GLU R7, GLUR 7, GluR 7a, GluR-7, GLUR7, GluR7a, Glutamate receptor 7, Glutamate receptor, Glutamate receptor ionotropic kainate 3, GRIK 3, GRIK3, GRIK3_HUMAN, ionotropic kainate 3.</p> <p>Background: Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration.</p> <p>Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neuro-transmission by glutamate, whereas the NMDA receptors are functionally characterized by a slow kinetic and a high permeability for Ca<sup>2+</sup> ions. The NMDA receptors consist of five subunits: epsilon 1, 2, 3, 4 and one zeta subunit. The zeta subunit is expressed throughout the brainstem, whereas the four epsilon subunits display limited distribution.</p>
Pathways:	<a href="#">Synaptic Membrane</a>

## Application Details

Application Notes:	<p>FCM 1:20-100</p> <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months