antibodies -online.com





anti-NSG2 antibody (AA 31-140) (Biotin)



Go to Froduct page

()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	NSG2 (HMP19)
Binding Specificity:	AA 31-140
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NSG2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NSG2/HMP19
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NSG2 (HMP19)
Alternative Name:	NSG2 (HMP19 Products)
Background: Synonyms: HMP19, Neuron-specic protein family member 2, Nsg2, NSG2_HUMAN, Protein family member 2,	

Background: With 181 million base pairs encoding around 1,000 genes, chromosome 5 is about 6 % of human genomic DNA. It is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5 associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome. Deletion of 5q or chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

Application Details

Restrictions:	For Research Use only
	IHC-F 1:100-500
Application Notes:	IHC-P 1:200-400

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months