



[Go to Product page](#)

Datasheet for ABIN1394854
anti-HTR1F antibody (AA 1-100) (Alexa Fluor 488)

Overview

Quantity:	100 µL
Target:	HTR1F
Binding Specificity:	AA 1-100
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HTR1F antibody is conjugated to Alexa Fluor 488
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human 5HT1F Receptor/SR-1F
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	HTR1F
Alternative Name:	5HT1F Receptor/SR-1F (HTR1F Products)

Target Details

Background: Synonyms: 5 HT 1F, 5 HT1F, 5 hydroxytryptamine serotonin receptor 1F, 5 hydroxytryptamine 1F receptor, 5 hydroxytryptamine receptor 1E beta, 5 hydroxytryptamine receptor 1F, 5 hydroxytryptamine serotonin receptor 1E beta, 5 hydroxytryptamine serotonin receptor 1F, 5HT 1F, 5HT1F, HTR 1F, HTR1EB, HTR1EL, HTR1F, MR 77, MR77, Serotonin receptor 1E beta, Serotonin receptor gene, 5HT1F_HUMAN.

Background: The members of the G-protein-coupled receptor family are distinguished by their slow transmitting response to ligand binding. These seven transmembrane proteins include the adrenergic, serotonin and dopamine receptors. The effect of the signaling molecule can be excitatory or inhibitory depending on the type of receptor to which it binds. b-adrenergic bound to adrenaline activates adenylyl cyclase, while a2-adrenergic receptor bound to adrenaline inhibits adenylyl cyclase. Like the a2-adrenergic receptor, serotonin receptor functions are also mediated by G proteins that inhibit the activity of adenylyl cyclase. The serotonin receptors have been classified into several categories, designated SR-17 (5HT17). Subtypes within the SR-1 group include SR-1A, -1B, -1D, -1E and -1F.

Pathways: [JAK-STAT Signaling](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months