

Datasheet for ABIN1394869

anti-Serotonin Receptor 2B antibody (AA 1-100) (Biotin)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Serotonin Receptor 2B (HTR2B)
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Serotonin Receptor 2B antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human 5HT2B Receptor/SR-2B
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	Serotonin Receptor 2B (HTR2B)
Alternative Name:	5HT2B Receptor/SR-2B (HTR2B Products)
Background:	Synonyms: 5 HT 2B, 5 HT 2B receptor, 5 HT2B, 5 HT2B, 5 hydroxytryptamine serotonin receptor

Target Details

2B, 5 hydroxytryptamine serotonin receptor 2B G protein coupled, 5 hydroxytryptamine 2B receptor, 5 hydroxytryptamine receptor 2B, 5-HT-2B, 5-HT2B, 5-hydroxytryptamine receptor 2B, 5HT2B, 5HT2B_HUMAN, 5HT2F, Htr 2b, Htr2b, NP75 protein, SEROTONIN 5HT2B RECEPTOR, Serotonin receptor 2B, Stomach fundus serotonin receptor.

Background: Serotonin (also designated 5-hydroxytryptamine or 5-HT) is a molecule that functions as a neurotransmitter, a hormone and a mitogen, and it is predominantly expressed in the gut, platelets and central nervous system (CNS). In the CNS, serotonin modulates several processes, including anxiety, sleep, appetite, behavior and drug abuse. In platelets and gut, serotonin plays a major role in cardiovascular function and motility of the gastrointestinal tract, respectively. Serotonin mediates its effects through several of G protein coupled receptors, designated 5-HT receptors or alternatively SR receptors. The SR-2 receptors are comprised of three subtypes, SR-2A, SR-2B and SR-2C, which activate phospholipase C and release intracellular stores of calcium in response to serotonin. SR-2A has a specific role in tracheal smooth muscle contraction, bronchoconstriction and mediating aldosterone production, and it is also thought to play a role in several psychiatric disorders, including depression and schizophrenia. SR-2B is expressed in embryonic and adult cardiovascular tissues, gut and brain and plays an important role in the pathology of cardiac disorders. SR-2C is thought to mediate the effects of atypical antipsychotic drugs.

Pathways: [JAK-STAT Signaling](#), [Inositol Metabolic Process](#), [Regulation of G-Protein Coupled Receptor Protein Signaling](#), [Regulation of Carbohydrate Metabolic Process](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months