



[Go to Product page](#)

Datasheet for ABIN1394905
anti-HTR7 antibody (AA 51-150) (Biotin)

Overview

Quantity:	100 µL
Target:	HTR7
Binding Specificity:	AA 51-150
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HTR7 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human 5HT7 Receptor/SR-7
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	HTR7
Alternative Name:	5HT7 Receptor/SR-7 (HTR7 Products)

Target Details

Background: Synonyms: HT 7, 5 HT X, 5 hydroxytryptamine serotonin receptor 7 adenylate cyclase coupled, 5 hydroxytryptamine serotonin receptor 7, 5 hydroxytryptamine receptor 7, 5-HT-7, 5-HT-X, 5-HT7, 5-hydroxytryptamine receptor 7, 5HT7, 5HT7R_HUMAN, HTR7, Serotonin 5 HT 7 receptor, Serotonin receptor 7.

Background: Serotonin (also designated 5-hydroxytryptamine or 5-HT) is a molecule that functions as a neurotransmitter, a hormone and a mitogen, and it is predominantly expressed in the gut, platelets and central nervous system (CNS). In the CNS, serotonin modulates several processes, including anxiety, sleep, appetite, behavior and drug abuse. In platelets and gut, serotonin plays a major role in cardiovascular function and motility of the gastrointestinal tract, respectively. Serotonin mediates its effects through several of G protein coupled receptors, designated 5-HT receptors or alternatively SR receptors. SR-3 is a ligand-gated ion channel, whereas all other known serotonin receptor subtypes are G protein-coupled receptors. SR-4 mediates widespread effects in central and peripheral nervous systems. SR-7 belongs to the superfamily of G protein-coupled receptors. The gene which encodes SR-7 maps to human chromosome 10q21-q24.

Gene ID: 3363

UniProt: [P34969](#)

Pathways: [JAK-STAT Signaling](#)

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months