



[Go to Product page](#)

Datasheet for ABIN1394947
anti-GABRB1 antibody (pSer434) (Biotin)

Overview

Quantity:	100 µL
Target:	GABRB1
Binding Specificity:	pSer434
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GABRB1 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic phosphopeptide derived from human GARB1 around the phosphorylation site of Ser434
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	GABRB1
---------	--------

Target Details

Alternative Name: GARB1 ([GABRB1 Products](#))

Background: Synonyms: GABA A Receptor beta 1 phospho S434; GARB1 Ser 434; GABAA receptor subunit beta-1; GABA-A receptor, beta-1 polypeptide; Gabrb-1; Gamma Aminobutyric Acid A Receptor Beta 1; Gamma Aminobutyric Acid Receptor , beta-1; Gamma-aminobutyric acid GABA A receptor, subunit beta 1; Gamma-aminobutyric acid receptor subunit beta-1; GARB1; GABRA1; AW061132; B230208N19Rik; GABAA receptor beta 1; GABAA receptor subunit beta-1; GABA-A receptor, beta-1 polypeptide; Gabrb-1; GABRB1; Gamma aminobutyric acid GABA A receptor beta 1; Gamma Aminobutyric Acid A Receptor Beta 1; Gamma Aminobutyric Acid Receptor , beta-1; Gamma-aminobutyric acid GABA A receptor, subunit beta 1; Gamma-aminobutyric acid receptor subunit beta-1; GARB1; GBRB1_HUMAN.

Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (g-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl⁻ conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R alpha 1-6, GABAA R beta 1-3, GABAA R ρ 1-3, GABAA R δ , GABAA R gamma, GABAA R delta 1 and GABAA R delta 2. The GABAB family is composed of GABAB R1 alpha and GABAB R1 beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2 and -3). The GABA transporters function to terminate GABA action.

Application Details

Application Notes: WB 1:300-5000
IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months