# antibodies -online.com





## anti-GABRB2 antibody (AA 251-350) (Biotin)



( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	GABRB2	
Binding Specificity:	AA 251-350	
Reactivity:	Rat, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GABRB2 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GABRB2/GABA A Receptor beta 2	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	

### **Target Details**

Target:	GABRB2
Alternative Name:	GABRB2/GABA A Receptor beta 2 (GABRB2 Products)

#### **Target Details**

Storage Comment:

Expiry Date:

rarget Details		
Background:	Synonyms: GABA, GABAA receptor subunit beta-2, GABRB2, Gamma aminobutyric acid GABA A	
	receptor, beta 2, Gamma aminobutyric acid A receptor beta 2, Gamma-aminobutyric acid	
	receptor subunit beta-2, Gamma-aminobutyric-acid receptor subunit beta-2, GBRB2_HUMAN,	
	Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the	
	production of GABA (gamma-aminobutyric acid). In the central nervous system GABA functions	
	as the main inhibitory transmitter by increasing a Cl- conductance that inhibits neuronal firing.	
	GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB)	
	receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are	
	ligand-gated ion channels, however, they are structurally and functionally distinct.	
Pathways:	Sensory Perception of Sound, Synaptic Membrane	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Store at -20°C for 12 months.

12 months