

[Go to Product page](#)

Datasheet for ABIN1395049

anti-GABRE antibody (AA 54-150) (Biotin)

Overview

Quantity:	100 µL
Target:	GABRE
Binding Specificity:	AA 54-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GABRE antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GABARE/GABA Receptor Epsilon
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Horse
Purification:	Purified by Protein A.

Target Details

Target:	GABRE
Alternative Name:	GABARE/GABA Receptor Epsilon (GABRE Products)
Background:	Synonyms: GABAA receptor epsilon, GABAA receptor subunit epsilon, Gabre, Gamma

Target Details

aminobutyric acid GABA A receptor epsilon, Gamma aminobutyric acid receptor subunit epsilon, Gamma-aminobutyric acid receptor subunit epsilon, GBRE_HUMAN.

Background: GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (gamma-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl⁻ conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA R Alpha 1-6, GABAA R Beta 1-3, GABAA R Gamma 1-3, GABAA R Delta, GABAA R Epsilon, GABAA R Zeta 1 and GABAA R Zeta 2. The GABAB family is composed of GABAB R1 Alpha and GABAB R1 Beta. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

Pathways: [Synaptic Membrane](#)

Application Details

Application Notes: IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Expiry Date: 12 months