
# antibodies -online.com





# anti-MPDZ antibody (AA 388-480) (Alexa Fluor 350)



Go to Product page

( )	۱ ۱	$\cap$	r	/1	$\cap$	۱ ۸	1
0	'V	ㄷ	ı١	νı	$\Box$	V	۷

Quantity:	100 μL
Target:	MPDZ
Binding Specificity:	AA 388-480
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MPDZ antibody is conjugated to Alexa Fluor 350
Application:	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MUPP1/MPDZ
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

## **Target Details**

Target:	MPDZ
Alternative Name:	MUPP1/MPDZ (MPDZ Products)

#### Target Details

Bac	kar	'nΙ	ın	Ч.
Duo		$\sim$	<i>.</i>	ч.

Synonyms: MPDZ, Multi PDZ domain protein 1, Multi-PDZ domain protein 1, multiple PDZ domain protein, MPDZ\_HUMAN.

Background: MPDZ is a 2,042 amino acid peripheral membrane protein that colocalizes with SR-2C on the apical membrane of epithelial choroid plexus cells. Expressed in heart, brain, placenta, liver, skeletal muscle, kidney and pancreas, MPDZ causes clustering of SR-2C, a serotonin receptor, at the cell surface. MPDZ is member of the NMDAR signaling complex that is involved in regulating AMPAR potentiation and synaptic plasticity in excitatory synapses. As a tight junction protein in epithelial cells, MPDZ interacts with SSTR3, a G-protein-coupled receptor, to regulate transepithelial permeability in a pertussis toxin sensitive manner. MPDZ, along with Kir4.2, may form a complex with other proteins in the nephron and regulate ion transport. MPDZ contains one L27 domain and thirteen PDZ domains.

Pathways:

Synaptic Membrane, Phototransduction

### **Application Details**

Application Notes:

FCM 1:20-100

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months