antibodies

## Datasheet for ABIN1395286 anti-SCN2B antibody (AA 65-180) (Alexa Fluor 488)



Overview

| Quantity:            | 100 µL  |
|----------------------|---|
| Target:              | SCN2B   |
| Binding Specificity: | AA 65-180   |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This SCN2B antibody is conjugated to Alexa Fluor 488  |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

## Product Details

| Immunogen:            | KLH conjugated synthetic peptide derived from human SCN2B |
|-----------------------|---|
| Isotype:              | lgG   |
| Predicted Reactivity: | Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Rabbit                  |
| Purification:         | Purified by Protein A.                                    |

## Target Details

| Target:           | SCN2B  |
|-------------------|--|
| Alternative Name: | SCN2B (SCN2B Products)   |
| Background:       | Synonyms: Neuronal voltage gated sodium channel beta 2 subunit, Scn 2b, Scn2b, |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1395286 | 03/07/2024 | Copyright antibodies-online. All rights reserved. SCN2B\_HUMAN, Sodium channel beta 2 subunit, Sodium channel subunit beta 2, Sodium channel subunit beta-2, Sodium channel voltage gated type II beta, Sodium channel voltage gated type II beta polypeptide.

Background: Voltage-gated sodium channels are selective ion channels that regulate the permeability of sodium ions in excitable cells. During the propagation of an action potential, sodium channels allow an influx of sodium ions, which rapidly depolarizes the cell. Na+ CP type II beta(sodium channel, voltage-gated, type II, beta), also known as SCN2B, is a 215 amino acid single-pass type I membrane protein that plays a critical role in the expression and assembly of the heterotrimeric complex of the sodium channel and interacts with Tenascin-R to influence the clustering and regulation of sodium channels at nodes of Ranvier. Expressed specifically in brain, Na+ CP type II beta contains one Ig-like C2-type (immunoglobulin-like) domain and is encoded by a gene that maps to human chromosome 11q23.3 and mouse chromosome 9 A5.2.

## Application Details

| Application Notes: | IF(IHC-P) 1:50-200   |
|--------------------|--|
|                    | IF(IHC-F) 1:50-200   |
|                    | IF(ICC) 1:50-200   |
| Restrictions:      | For Research Use only  |
| Handling           |  |
| Format:            | Liquid   |
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.        |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                                  |
| Expiry Date:       | 12 months  |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1395286 | 03/07/2024 | Copyright antibodies-online. All rights reserved.