

[Go to Product page](#)

Datasheet for ABIN1395298

anti-SLC6A7 antibody (AA 151-260) (Alexa Fluor 488)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | SLC6A7 |
| Binding Specificity: | AA 151-260 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This SLC6A7 antibody is conjugated to Alexa Fluor 488 |
| Application: | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| | |
|-----------------------|---|
| Immunogen: | KLH conjugated synthetic peptide derived from human SLC6A7/PROT |
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|---|
| Target: | SLC6A7 |
| Alternative Name: | SLC6A7/PROT (SLC6A7 Products) |
| Background: | Synonyms: SC6A7_HUMAN, Slc6a7, Sodium-dependent proline transporter, Solute carrier |

Target Details

family 6 member 7.

Background: The GAT1 gene family includes sodium- and chloride-dependent plasma membrane transporters for neurotransmitters, metabolites and osmolites, which couple substrate flux to transmembrane electrochemical gradients. PROT (Sodium-dependent proline transporter), also known as Solute carrier family 6 member 7, is a 636 amino acid multi-pass membrane protein that is a GAT1 family member specifically expressed in regions of the brain. PROT terminates the action of proline by its high affinity sodium/chloride-dependent reuptake into pre-synaptic terminals. Enriched in glutamatergic synaptic terminals, it is likely that PROT plays an important role in excitatory events of neurotransmission. PROT-mediated proline uptake is inhibited by compounds such as benztropine, LP-403812 and Des-Tyr-Leu-enkephalin (GGFL). These inhibitors of proline uptake may lead to the development of therapeutic agents for certain neurologic disorders.

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months