

[Go to Product page](#)

Datasheet for ABIN1395364

anti-STRN3 antibody (AA 75-180) (Alexa Fluor 488)

Overview

Quantity:	100 µL
Target:	STRN3
Binding Specificity:	AA 75-180
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STRN3 antibody is conjugated to Alexa Fluor 488
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Striatin 3
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	STRN3
Alternative Name:	Striatin 3/SG2NA (STRN3 Products)

Target Details

Background:	<p>Synonyms: Striatin3, Striatin-3, Cell cycle autoantigen SG2NA, Cell cycle S/G2 nuclear autoantigen, GS2NA, nuclear autoantigen, S/G2 antigen, SG2NA, Striatin 3, Striatin, calmodulin binding protein 3, STRN3_HUMAN.</p> <p>Background: Striatin, SG2NA, and zinedin, the three mammalian members of the striatin family, are multimodular, WD repeat and calmodulin-binding proteins. Zinedin and SG2NA share with striatin identical protein-protein interaction domains and the same overall domain structure. All three proteins are both cytosolic and membrane-bound and bind calmodulin in the presence of calcium. Striatin is a neuronal, intracellular protein strictly expressed in the somato-dendritic compartment, including spines and subsets of neurons, and is considered as a marker of neuronal polarity. Downregulation of striatin, which is expressed in a few subsets of neurons, impairs the growth of dendrites as well as rat locomotor activity. Zinedin is mainly expressed in the central nervous system, whereas SG2NA is mainly expressed in the brain and muscle.</p>
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway , Regulation of Intracellular Steroid Hormone Receptor Signaling

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months