

[Go to Product page](#)

Datasheet for ABIN1395389

anti-DLGAP2 antibody (AA 251-360) (Alexa Fluor 555)

Overview

Quantity:	100 µL
Target:	DLGAP2
Binding Specificity:	AA 251-360
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DLGAP2 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SAPAP2/DLGAP2
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse
Purification:	Purified by Protein A.

Target Details

Target:	DLGAP2
Alternative Name:	SAPAP2/DLGAP2 (DLGAP2 Products)

Target Details

Background:	<p>Synonyms: DAP-2, DAP2, Discs large Drosophila homolog associated protein 2, Discs large homolog associated protein 2, Disks large-associated protein 2, Dlgap2, DLGP2_HUMAN, PSD-95/SAP90-binding protein 2, SAP90/PSD-95-associated protein 2, SAPAP2, hide.</p> <p>Background: A guanylate kinase is a phosphotransferase that produces ADP and GDP from the substrates ATP and GMP. SAPAP2, also known as DAP-2 (Disks large-associated protein 2) and PSD-95/SAP90-binding protein 2, is a 1054 amino acid protein that localizes to the postsynaptic membrane of neuronal cells of the brain and kidney. SAPAP2 likely acts as a signaling molecule which interacts with the human genes DLG1 and DLG4/PSD-95. The gene encoding SAPAP2, DLGAP2, maps to human chromosome 8. Consisting of nearly 146 million base pairs, chromosome 8 encodes over 800 genes and is associated with a variety of diseases and malignancies. Schizophrenia, bipolar disorder, Trisomy 8, Pfeiffer syndrome, congenital hypothyroidism, Waardenburg syndrome and some leukemias and lymphomas are thought to occur as a result of defects in specific genes that maps to chromosome 8.</p>
-------------	---

Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months