antibodies -online.com





Datasheet for ABIN1395428

anti-CARTPT antibody (AA 31-116) (FITC)



()	1 /	0	rv	/ 1 /	71	Α.
	1//	\vdash	1 \/	16		1/1/
\sim	v	\sim	1 V	١,	_	v v

Quantity:	100 μL	
Target:	CARTPT	
Binding Specificity:	AA 31-116	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CARTPT antibody is conjugated to FITC	
Application:	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CART	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	CARTPT	
Alternative Name:	CART (CARTPT Products)	
Background:	Synonyms: CART, CART prepropeptide, Cocaine and amphetamine regulated transcript,	

CART_HUMAN.

Background: The CART gene encodes for a protein which has an important role in the regulation of appetite and body weight. The CART (cocaine- and amphetamine-regulated transcript) neuropeptide is an mRNA that changes in response to psychostimulant drug administration. Injection of CART peptides into the ventral tegmental area produces psychostimulant-like effects. CART localizes to areas of the central and peripheral nervous systems and is involved in feeding behavior when injected centrally. Expression of CART in the rat hypothalamus is modulated by nutritional status, and injection of synthetic CART peptide into the forebrain ventricular system suppresses food intake, indicating a possible role in hypothalamic control of energy homeostasis. Its identification in cell bodies and central terminals of vagal afferent neurons additionally suggests a role in brainstem mechanisms of meal termination and satiety.

Pathways:

Hormone Transport, Negative Regulation of Hormone Secretion, Carbohydrate Homeostasis, Feeding Behaviour

Application Details

Application Notes: IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months