# antibodies -online.com





## anti-KCNT2 antibody (AA 51-150) (Alexa Fluor 647)



_					
U	V	er	V	Ie	W

Quantity:	100 μL	
Target:	KCNT2	
Binding Specificity:	AA 51-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This KCNT2 antibody is conjugated to Alexa Fluor 647	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human KCNT2
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Chicken, Rabbit
Purification:	Purified by Protein A.

### **Target Details**

Target:	KCNT2
Alternative Name:	KCNT2 (KCNT2 Products)
Background: Synonyms: KCa4.2, Potassium channel, subfamily T, member 2, Sequence like an inte	

conductance potassium channel subunit, SLICK, SLO2.1, Sodium and chloride activated ATP sensitive potassium channel, Sodium and chloride activated ATP sensitive potassium channel Slo2.1, KCNT2\_HUMAN.

Background: Voltage-gated K+ channels in the plasma membrane are important regulators of electrical signaling, controlling the repolarization and the frequency of action potentials in neurons, muscles and other excitable cells. KCNT2 is a 1,135 amino acid multi-pass transmembrane protein belonging to the potassium channel family (calcium-activated subfamily) of proteins. KCNT2 produces rapidly activating outward rectifier potassium currents in reponse to high intracellular sodium and chloride levels. Its channel activity is inhibited by ATP, inhalation anesthetics, such as isoflourane, and upon stimulation of G-protein coupled receptors, such as mAChR M1 and GluR-1. There are four isoforms of KCNT2 that are produced as a result of alternative splicing events.

### **Application Details**

Restrictions:	For Research Use only
	IF(ICC) 1:50-200
	IF(IHC-F) 1:50-200
Application Notes:	IF(IHC-P) 1:50-200

#### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	