antibodies -online.com





anti-Kv2.2 antibody (AA 21-120) (Alexa Fluor 555)



Go to Product page

| () | 11/0 | K\ / | iew |
|-----|--------|--------|-------------|
| | \cup | 'I V/I | $I \cap VV$ |
| | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | Kv2.2 (KCNB2) |
| Binding Specificity: | AA 21-120 |
| Reactivity: | Rat, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Kv2.2 antibody is conjugated to Alexa Fluor 555 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human Kv2.2 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Predicted Reactivity: | Human,Dog,Sheep,Pig,Horse,Chicken,Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | Kv2.2 (KCNB2) |
|-------------------|------------------------|
| Alternative Name: | Kv2.2 (KCNB2 Products) |

Target Details

Background:

Synonyms: delayed rectier potassium channel protein, KCNB2, KCNB2_HUMAN, potassium channel Kv2.2, potassium voltage gated channel subfamily B member 2, Potassium voltage-gated channel subfamily B member 2, Voltage-gated potassium channel subunit Kv2.2.

Background: Voltage-gated K+ channels in the plasma membrane control the repolarization and the frequency of action potentials in neurons, muscles and other excitable cells. The KV gene family encodes more than 30 proteins that comprise the subunits of the K+ channels, and they vary in their gating and permeation properties, subcellular distribution and expression patterns. Functional KV channels assemble as tetramers consisting of pore-forming alpha subunits (KV), which include the KV1, KV2, KV3, KV4 and KV9 proteins, and accessory or KV-subunits that modify the gating properties of the coexpressed KV subunits. KV2.2 is a multi-pass membrane protein that regulates the voltage-dependent K+ permeability of excitable membranes. Its tail may be influential in the targeting of the channel to specific subcellular compartments and/or the regulation of channel activity.

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|-----------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |