

Datasheet for ABIN1396566

anti-KCNMA1 antibody (AA 21-120) (Biotin)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	KCNMA1
Binding Specificity:	AA 21-120
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KCNMA1 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BKCA alpha
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	KCNMA1
Alternative Name:	BK channel (KCNMA1 Products)

Target Details

Background:	<p>Synonyms: subfamily M subunit alpha-1, BK channel, BKCA alpha, BKCA alpha subunit, BKTm, Calcium-activated potassium channel, Calcium-activated potassium channel subunit alpha-1, Drosophila slowpoke like, hSlo, KVCAalpha, KCa1.1, KCMA1_HUMAN, KCNMA, KCNMA1, Maxi K channel, Maxi Potassium channel alpha, MaxiK, SAKCA, SLO alpha, SLO, Slo homolog, Slo-alpha, Slo1, Slowpoke homolog.</p> <p>Background: MaxiK channels are large conductance, voltage and calcium-sensitive potassium channels which are fundamental to the control of smooth muscle tone and neuronal excitability. MaxiK channels can be formed by 2 subunits: the pore-forming alpha subunit, which is the product of this gene, and the modulatory beta subunit. Intracellular calcium regulates the physical association between the alpha and beta subunits. Alternatively spliced transcript variants encoding different isoforms have been identified. [provided by RefSeq, Jul 2008].</p>
Pathways:	Regulation of Hormone Metabolic Process , Sensory Perception of Sound

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months