

Datasheet for ABIN1397000

## anti-TGFB1 antibody (AA 101-200) (AbBy Fluor® 350)



[Go to Product page](#)

### Overview

|                      |                                                                                                                                   |
|----------------------|-----------------------------------------------------------------------------------------------------------------------------------|
| Quantity:            | 100 µL                                                                                                                            |
| Target:              | TGFB1                                                                                                                             |
| Binding Specificity: | AA 101-200                                                                                                                        |
| Reactivity:          | Human, Mouse                                                                                                                      |
| Host:                | Rabbit                                                                                                                            |
| Clonality:           | Polyclonal                                                                                                                        |
| Conjugate:           | This TGFB1 antibody is conjugated to AbBy Fluor® 350                                                                              |
| Application:         | Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)),<br>Immunofluorescence (Cultured Cells) (IF (cc)) |

### Product Details

|                   |                                                                                                                                                    |
|-------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen:        | KLH conjugated synthetic peptide derived from human TGF beta 1/LAP                                                                                 |
| Isotype:          | IgG                                                                                                                                                |
| Specificity:      | The antibody targets the Latency-associated peptide (LAP), which is a cleavage product of the Transforming growth factor beta-1 proprotein (TGFB1) |
| Cross-Reactivity: | Human, Mouse                                                                                                                                       |
| Purification:     | Purified by Protein A.                                                                                                                             |

### Target Details

|         |       |
|---------|-------|
| Target: | TGFB1 |
|---------|-------|

## Target Details

|                   |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
|-------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Alternative Name: | TGFB1 ( <a href="#">TGFB1 Products</a> )                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| Background:       | <p>Synonyms: Transforming growth factor beta-1 proprotein, CED, LAP, DPD1, TGFB, TGFbeta, TGF-beta-1, TGFB1</p> <p>Background: Latency-associated peptide: Required to maintain the Transforming growth factor beta-1 (TGF-beta-1) chain in a latent state during storage in extracellular matrix. Associates non-covalently with TGF-beta-1 and regulates its activation via interaction with 'milieu molecules', such as LTBP1, LRRC32/GARP and LRRC33/NRROS, that control activation of TGF-beta-1. Interaction with LRRC33/NRROS regulates activation of TGF-beta-1 in macrophages and microglia (Probable). Interaction with LRRC32/GARP controls activation of TGF-beta-1 on the surface of activated regulatory T-cells (Tregs). Interaction with integrins (ITGAV:ITGB6 or ITGAV:ITGB8) results in distortion of the Latency-associated peptide chain and subsequent release of the active TGF-beta-1.</p> |
| Gene ID:          | 7040                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
| UniProt:          | <a href="#">P01137</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Pathways:         | <a href="#">EGFR Signaling Pathway</a> , <a href="#">Dopaminergic Neurogenesis</a> , <a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Glycosaminoglycan Metabolic Process</a> , <a href="#">Regulation of Leukocyte Mediated Immunity</a> , <a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Positive Regulation of Immune Effector Process</a> , <a href="#">Cell-Cell Junction Organization</a> , <a href="#">Production of Molecular Mediator of Immune Response</a> , <a href="#">Ribonucleoside Biosynthetic Process</a> , <a href="#">Skeletal Muscle Fiber Development</a> , <a href="#">Regulation of Carbohydrate Metabolic Process</a> , <a href="#">Protein targeting to Nucleus</a> , <a href="#">Autophagy</a> , <a href="#">Cancer Immune Checkpoints</a>                                                                                      |

## Application Details

|                    |                                                              |
|--------------------|--------------------------------------------------------------|
| Application Notes: | IF(IHC-P) 1:50-200<br>IF(IHC-F) 1:50-200<br>IF(ICC) 1:50-200 |
| Restrictions:      | For Research Use only                                        |

## Handling

|                |                                                                                              |
|----------------|----------------------------------------------------------------------------------------------|
| Format:        | Liquid                                                                                       |
| Concentration: | 1 µg/µL                                                                                      |
| Buffer:        | Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |

## Handling

---

50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months