antibodies .- online.com







anti-CRTAM antibody (AA 31-130) (Alexa Fluor 488)



()	11/0	K\ /	iew	1
	\cup	ועוי	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	CRTAM	
Binding Specificity:	AA 31-130	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CRTAM antibody is conjugated to Alexa Fluor 488	
Application:	Flow Cytometry (FACS), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CRTAM
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	CRTAM
Alternative Name:	CRTAM (CRTAM Products)

Target Details

Our arrange ODOFF Outstands and regulation Toolly and a little outside LT
Synonyms: CD355, Cytotoxic and regulatory T-cell molecule, Class-I MHC-restricted T-cell-
associated molecule, CRTAM
Background: Interaction with CADM1 promotes natural killer (NK) cell cytotoxicity and
interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated
rejection of tumors expressing CADM3 in vivo.
56253
095727
Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process,
Activated T Cell Proliferation
FCM 1:20-100
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Limid
Liquid
1 μg/μL
Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months