Datasheet for ABIN1397433 anti-STXBP3 antibody (AA 165-260) (Biotin)

antibodies.com



Overview

Quantity:	100 μL	
Target:	STXBP3	
Binding Specificity:	AA 165-260	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This STXBP3 antibody is conjugated to Biotin	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Unc18-3
Isotype:	lgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Pig,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	STXBP3
Alternative Name:	Unc18-3 (STXBP3 Products)
Background:	Synonyms: Munc 18 3, Munc18 3, Munc 18-3, Protein unc-18 homolog 3, STXB3_HUMAN.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1397433 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

## Target Details

Background: May participate in the regulation of synaptic vesicle docking through interaction with GTP-binding proteins. Essential for neurotransm syntaxin, a component of the synaptic vesicle fusion machinery probably interact with syntaxins 1, 2, and 3 but not syntaxin 4. May play a role in despecificity of intracellular fusion reactions.         Gene ID:       6814         Pathways:       Hormone Transport         Application Details       IHC-P 1:200-400         IHC-F 1:100-500       IHC-F 1:100-500         Restrictions:       For Research Use only         Handling       Iuquid         Concentration:       1 µg/µL         Buffer:       Aqueous buffered solution containing 0.01M TBS (pH 7,4) with 1 % BSA, 50 % Glycerol.         Preservative:       ProClin         Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C         Storage Comment:       Storage 2:0°C for 12 months.	
syntaxin, a component of the synaptic vesicle fusion machinery probably interact with syntaxins 1, 2, and 3 but not syntaxin 4. May play a role in de specificity of intracellular fusion reactions.Gene ID:6814Pathways:Hormone TransportApplication DetailsIHC-P 1:200-400 IHC-F 1:100-500Restrictions:For Research Use onlyHandlingIuquidFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.	ing and fusion, possibly
interact with syntaxins 1, 2, and 3 but not syntaxin 4. May play a role in despecificity of intracellular fusion reactions. Gene ID: 6814 Pathways: Hormone Transport Application Details IHC-P 1:200-400 IHC-F 1:100-500 Restrictions: For Research Use only Handling Format: Liquid Concentration: 1 µg/µL Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol. Preservative: ProClin Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only. Storage: -20 °C	smission and binds
specificity of intracellular fusion reactions.         Gene ID:       6814         Pathways:       Hormone Transport         Application Details       IHC-P 1:200-400         Application Notes:       IHC-P 1:200-400         IHC-F 1:100-500       IHC-F 1:100-500         Restrictions:       For Research Use only         Handling       Iuquid         Concentration:       1 µg/µL.         Buffer:       Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.         Preservative:       ProClin         Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C	bly in a 1:1 ratio. Can
Gene ID:       6814         Pathways:       Hormone Transport         Application Details       IHC-P 1:200-400         Application Notes:       IHC-P 1:200-400         IHC-F 1:100-500       IHC-F 1:100-500         Restrictions:       For Research Use only         Handling       Iuguid         Concentration:       1 µg/µL         Buffer:       Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.         Preservative:       ProClin         Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C	determining the
Pathways:       Hormone Transport         Application Details         Application Notes:       IHC-P 1:200-400 IHC-F 1:100-500         Restrictions:       For Research Use only         Handling         Format:       Liquid         Concentration:       1 µg/µL         Buffer:       Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.         Preservative:       ProClin         Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C	
Application Details         Application Notes:       IHC-P 1:200-400 IHC-F 1:100-500         Restrictions:       For Research Use only         Handling       Iuquid         Format:       Liquid         Concentration:       1 µg/µL         Buffer:       Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.         Preservative:       ProClin         Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C	
Application Notes:IHC-F 1:200-400 IHC-F 1:100-500Restrictions:For Research Use onlyHandlingIuquidFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
IHC-F 1:100-500Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
Restrictions:For Research Use onlyHandlingFormat:LiquidConcentration:1 µg/µLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
HandlingFormat:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
Format:LiquidConcentration:1 μg/μLBuffer:Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
Concentration:1 µg/µLBuffer:Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
Buffer:Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 50 % Glycerol.Preservative:ProClinPrecaution of Use:This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.Storage:-20 °C	
50 % Glycerol.         Preservative:       ProClin         Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C	
Precaution of Use:       This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTA handled by trained staff only.         Storage:       -20 °C	SA, 0.03 % Proclin300 and
handled by trained staff only.       Storage:	
Storage: -20 °C	TANCE, which should be
Storage Comment: Store at -20°C for 12 months.	
Expiry Date: 12 months	