

[Go to Product page](#)

Datasheet for ABIN1397479

**anti-ITPR3 antibody (AA 21-120) (Alexa Fluor 555)**

## Overview

Quantity:	100 µL
Target:	ITPR3
Binding Specificity:	AA 21-120
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ITPR3 antibody is conjugated to Alexa Fluor 555
Application:	Western Blotting (WB), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ITPR3
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	ITPR3
Alternative Name:	Itpr3 ( <a href="#">ITPR3 Products</a> )

## Target Details

Background:	<p>Synonyms: IP3R-III, IP3 receptor isoform 3, 4 antibody 5-trisphosphate receptor, 5-trisphosphate receptor type 3, FLJ36205, Inositol 1, Inositol 1,4,5 trisphosphate receptor type 3, IP3 receptor, IP3R 3, IP3R, IP3R3, ITPR 3, ITPR3, ITPR3_HUMAN, Type 3 inositol 1, Type 3 inositol 1,4,5 trisphosphate receptor, Type 3 InsP3 receptor.</p> <p>Background: Inositol 1,4,5-trisphosphate (IP3) functions as a second messenger for a myriad of extracellular stimuli including hormones, growth factors and neurotransmitters. Receptor tyrosine kinases indirectly increase the intracellular levels of IP3 through the activation of phospholipases such as phospholipase C (PLC), which convert phosphatidylinositol-4,5 biphosphate into IP3 and diacylglycerol (DAG). The inositol 1,4,5-trisphosphate receptor, IP3R, acts as an inositol triphosphate (IP3)-gated calcium release channel in a variety of cell types. Three IP3 receptor subtypes have been described and are designated IP3R-I, IP3R-II and IP3R-III. IP3R-I is the predominant IP3R subtype expressed in neuronal tissues and the central nervous system, but is also expressed at high levels in the liver.</p>
Gene ID:	3710
Pathways:	<a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">EGFR Signaling Pathway</a> , <a href="#">Neurotrophin Signaling Pathway</a> , <a href="#">Thyroid Hormone Synthesis</a> , <a href="#">Myometrial Relaxation and Contraction</a> , <a href="#">G-protein mediated Events</a> , <a href="#">Interaction of EGFR with phospholipase C-gamma</a> , <a href="#">BCR Signaling</a>

## Application Details

Application Notes:	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

---

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months