## antibodies -online.com





## anti-AKAP5 antibody (AA 611-714) (AbBy Fluor® 647)



		do to i roduct page

Overview		
Quantity:	100 μL	
Target:	AKAP5	
Binding Specificity:	AA 611-714	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This AKAP5 antibody is conjugated to AbBy Fluor® 647	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human AKAP5	
Isotype:	IgG	
Cross-Reactivity:	Rat	
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	AKAP5	
Alternative Name:	AKAP5 (AKAP5 Products)	

## **Target Details**

Target Type:	Viral Protein	
Background:	Synonyms: A kinase PRKA anchor protein 5, A kinase anchor protein 5, A kinase anchor protein	
	79 kDa, A kinase anchoring protein 75/79, A-kinase anchor protein 5, A-kinase anchor protein 79	
	kDa, AKAP 5, AKAP 75, AKAP 79, AKAP-5, Akap5, AKAP5_HUMAN, AKAP75, AKAP79, cAMP	
	dependent protein kinase regulatory subunit II high affinity binding protein, cAMP-dependent	
	protein kinase regulatory subunit II high affinity-binding protein, H21.	
	Background: The type II cAMP-protein kinase (PKA) is a multifunctional kinase with a broad	
	range of substrates (1). Specificity of PKA signaling is thought to be mediated by the	
	compartmentalization of the kinase to specific sites within the cell. To maintain this specific	
	localization, the R subunit (RII) of PKA interacts with specific RII-anchoring proteins. This family	
	of proteins has been designated A-kinase anchoring proteins (AKAP) (1-3). Members of this	
	family, including MAP2 (microtubule-associated protein 2), neuronally expressed AKAP 79 and	
	AKAP 150, and the DNA binding AKAP 95, display differential tissue specificity and localization	
	(4-6). Evidence suggests that AKAP 79 and AKAP 150 are both capable of anchoring PKA to	
	postsynaptic densities (PSD), which are a network of proteins located on the internal surfaces	
	of excitatory synapses.	
Gene ID:	9495	
UniProt:	P24588	
Pathways:	cAMP Metabolic Process	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	