antibodies -online.com







anti-DCST1 antibody (AA 265-370) (AbBy Fluor® 555)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Quantity:	100 μL
Target:	DCST1
Binding Specificity:	AA 265-370
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DCST1 antibody is conjugated to AbBy Fluor® 555
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DCST1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	DCST1
Alternative Name:	DCST1 (DCST1 Products)

Target Details

Storage Comment:

rarget Details		
Background:	Synonyms: RP11 307C12.10, DC STAMP domain containing 1, FLJ32785, DC-STAMP domain-	
	containing protein 1, DCST1_HUMAN.	
	Background: Chromosome 1 is the largest human chromosome spanning about 260 million	
	base pairs and making up 8 % of the human genome. There are about 3,000 genes on	
	chromosome 1, and considering the great number of genes there are also a large number of	
	diseases associated with chromosome 1. Notably, the rare aging disease Hutchinson-Gilford	
	progeria is associated with the LMNA gene which encodes lamin A. When defective, the LMNA	
	gene product can build up in the nucleus and cause characteristic nuclear blebs. The	
	mechanism of rapidly enhanced aging is unclear and is a topic of continuing exploration. The	
	MUTYH gene is located on chromosome 1 and is partially responsible for familial adenomatous	
	polyposis. Stickler syndrome, Parkinsons, Gaucher disease and Usher syndrome are also	
	associated with chromosome 1. A breakpoint has been identified in 1q which disrupts the	
	DISC1 gene and is linked to schizophrenia. Aberrations in chromosome 1 are found in a variety	
	of cancers including head and neck cancer, malignant melanoma and multiple myeloma.	
Gene ID:	149095	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

		1.	
\vdash	land	lın	
	ıaııu	1111	U

Expiry Date:

12 months