

## Datasheet for ABIN1398314 anti-LGI4 antibody (AA 101-200) (FITC)



Overview Quantity: 100 µL LGI4 Target: Binding Specificity: AA 101-200 Reactivity: Human Rabbit Host: Clonality: Polyclonal Conjugate: This LGI4 antibody is conjugated to FITC Application: Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human LGI4
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	LGI4
Alternative Name:	LGI4 (LGI4 Products)
Background:	Synonyms: Leucine rich glioma inactivated gene 4, Leucine rich glioma inactivated protein 4,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN1398314 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

lauging righ report 1 OI family marches 4. Lauging righ report 1 OI family and 1 OI 1
leucine-rich repeat LGI family member 4, Leucine rich repeat LGI family member 4, LGI1 like protein 3, LGI4, Lgi4 protein, Lgil3, LGI4_HUMAN.
Background: LGI4, also known as leucine-rich glioma-inactivated protein 4, is a 537 amino acid
secreted glycosylated protein that is widely expressed, with highest levels found within the
nervous system. Interestingly, siRNA knockdown studies of LGI4 expression in Schwann cells
have been shown to result in the inhibition of myelination, thus suggesting that LGI4 is an
essential component of myelin formation and axon segregation. LGI4 shares significant
homology with its other family members, LGI1, LGI2 and LGI3. Significantly, mutations in the
gene encoding LGI1 have been linked to human temporal lobe epilepsy and, given the sequence
similarity of LGI4, it is likely that it also may be implicated in the pathology of seizures. LGI4 is
localized subcellularly to the Golgi, ER and vesicles. There are two isoforms of LGI4 that are
produced as a result of alternative splicing events.
163175
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
For Research Use only
Liquid
1 µg/µL
Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
50 % Glycerol.
ProClin
This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
handled by trained staff only.
-20 °C
Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/2 | Product datasheet for ABIN1398314 | 07/26/2024 | Copyright antibodies-online. All rights reserved.