antibodies - online.com







anti-RDH13 antibody (AA 101-200) (Alexa Fluor 488)



| \sim | | | | | |
|--------|----|----|----|-------------|---|
| () | VE | ۲۱ | /1 | \triangle | Λ |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | RDH13 | |
| Binding Specificity: | AA 101-200 | |
| Reactivity: | Mouse | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This RDH13 antibody is conjugated to Alexa Fluor 488 | |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human RDH13 | |
|-----------------------|---|--|
| Isotype: | IgG | |
| Cross-Reactivity: | Mouse | |
| Predicted Reactivity: | Human,Rat,Dog,Cow,Sheep,Horse,Rabbit | |
| Purification: | Purified by Protein A. | |

Target Details

| Target: | RDH13 |
|-------------------|------------------------|
| Alternative Name: | RDH13 (RDH13 Products) |

Target Details

| Background: | Synonyms: RDH13, RDH13_HUMAN, Retinol dehydrogenase 13 all trans and 9 cis, Retinol | | |
|---------------------|--|--|--|
| | dehydrogenase 13 all trans/9 cis, Retinol dehydrogenase 13. | | |
| | Background: RDH13, also known as all-trans and 9-cis retinol dehydrogenase 13 or SDR7C3, is | | |
| | a 331 amino acid mitochondrial protein belonging to the short-chain | | |
| | dehydrogenases/reductases (SDR) family. Widely expressed, mostly in eye, pancreas, placenta | | |
| | and lung, RDH13 localizes on the outer side of the inner mitochondrial membrane. Related to | | |
| | microsomal retinoid oxidoreductase RDH11, RDH13 is considered to be a major enzyme | | |
| | among the RDH family of proteins. Catalytically active, RDH13 recognizes retinoids as | | |
| | substrates and may function in retinoic acid production. RDH13 may function to protect the | | |
| | mitochondria against oxidative stress. Leber congenital amaurosis (LCA) type 3, an inherited | | |
| | autosomal recessive retinal disease, has been associated with defects of RDH13. LCA | | |
| | represents the most common genetic cause of congenital visual impairment in infants and | | |
| | children. | | |
| Gene ID: | 112724 | | |
| Application Details | | | |
| Application Notes: | IF(IHC-P) 1:50-200 | | |
| | IF(IHC-F) 1:50-200 | | |
| | IF(ICC) 1:50-200 | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 μg/μL | | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | | |
| | 50 % Glycerol. | | |
| Preservative: | ProClin | | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | | |
| | handled by trained staff only. | | |
| Storage: | -20 °C | | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | | |
| Expiry Date: | 12 months | | |
| | | | |